

विआई कैप्सुलर पॉलीसैकेराइड टाइफोयिड वैक्सीन आई.पी.
Vi Capsular Polysaccharide Typhoid Vaccine I.P.

TYPBAR®

टाईपबार®

Description

TYPBAR® is a sterile solution for intramuscular use containing the cell surface Vi polysaccharide extracted from Salmonella typhi Ty2 strain. The vaccine appears as a clear, colourless solution.

Composition

Each 0.5 ml dose contains :

Vi Capsular polysaccharide of Salmonella typhi (Ty2 strain) 25 µg, Phenol and buffer solution containing sodium chloride, disodium phosphate, monosodium phosphate and water for injection.

CLINICAL PARTICULARS

Therapeutic indications

TYPBAR® is indicated for active immunization against typhoid fever for both adults and children two years of age or older.

Selective immunization with **TYPBAR®** is recommended for the following :

- Travellers to high endemic areas.
- Household contacts of carriers.
- Healthcare personnel.
- Police, Armed forces and such other regimented personnel.

Contraindications

TYPBAR® should not be administered to subjects with known hypersensitivity to any component of the vaccine or to subjects having shown signs of hypersensitivity after a previous administration.

Dosage

The immunizing dose for adults and children 2 years of age and older is a single dose of 0.5 mL.

Subjects who remain at risk of typhoid fever should be given a single booster dose of the vaccine with an interval of not more than 3 years.

Method of administration

TYPBAR® is for intramuscular injection only. DO NOT inject intravenously.

TYPBAR® in adults should be given intramuscularly in the deltoid and children should be injected intramuscularly either in the deltoid or the vastus lateralis. **TYPBAR®** should not be injected into the gluteal area or areas where there may be a nerve trunk.

While using the multi-dose vial, care must be taken to use separate sterile syringes and needles for the administration of every dose.

Before use, **TYPBAR®** should be well shaken. Vaccine should be visually checked for the presence of any particulate matter or other coloration, if any, prior to its administration. If in doubt, do not use the contents of the vial.

Special Precautions and Warnings for Use

TYPBAR® protects against typhoid fever caused by Salmonella typhi. Protection is not conferred against Salmonella Paratyphi and other non-Typhoidal Salmonellae.

Adrenaline injection must be kept readily available following immunization, should an anaphylactic or other allergic reaction occur due to any component of the vaccine.

The administration of **TYPBAR®** should be delayed in subjects with acute infection or febrile illness.

TYPBAR® should be administered with caution to subjects with thrombocytopenia or bleeding disorders since bleeding may occur following an intramuscular administration in these subjects. It may be expected that in patients receiving immunosuppressive treatment or patients with immunodeficiency, an adequate response may not be achieved.

TYPBAR® should not be mixed with other vaccines or medicinal products in the same syringe.

Pregnancy

The effect of the **TYPBAR®** on fetal development or reproduction capacity has not been evaluated.

TYPBAR® should be given to a pregnant woman only if clearly needed.

Nursing Mothers

It is not known if **TYPBAR®** is excreted in human milk. It may be administered to nursing mothers only if clearly needed.

Paediatric Use

Safety and effectiveness of **TYPBAR®** in children 2 years of age and below has not been established. Polysaccharide vaccines in general have lower immunogenicity under this age.

ADVERSE REACTIONS

Most recipients of typhoid vaccine experience some reactions upon vaccination. These are generally moderate and short in duration. They mainly consist of local reactions at the injection site (erythema, induration and tenderness). Systemic reactions (malaise, headache, diarrhoea, vomiting myalgia and elevated temperature) are reported less commonly.

In very rare cases, allergic - type reactions (pruritus, rash urticaria) may be observed.

PHARMACEUTICAL PARTICULARS:

Category	:	Active immunizing agent.
Pharmaceutical form	:	Liquid for injection.
Shelf life	:	3 years from the date of manufacture.
Storage	:	at 2°C to 8°C.

DO NOT FREEZE. DISCARD IF FROZEN.

KEEP OUT OF REACH OF CHILDREN.

PRESENTATION:

TYPBAR® is presented in USP type 1 glass vial and PFS		
Single dose	:	0.5 mL.
Multi-dose	:	2.5 mL.
Multi-dose	:	5.0 mL.
Single Dose PFS	:	0.5 mL.

For additional information please write to:
Bharat Biotech International Limited
Shameerpet, Hyderabad-500 078. India.

Manufactured & Marketed by
 **Bharat Biotech**
INTERNATIONAL LIMITED
Genome Valley, Shameerpet,
Hyderabad - 500 078. India.

Warning: To be sold by
retail on the prescription
of a Registered Medical
Practitioner only